

Acrosyndromes

En médecine interne

Agglutinine froide





sclérodémie



sclérodermie





dermatomyosite




dermatomyosite

mégacapillaires









ROLLES

00 00200



A close-up photograph of the plantar surface of a foot, focusing on the second and third toes. The skin around the nail plates of both toes shows signs of inflammation, including redness and swelling, which is characteristic of subacute paronychia (SAPL). The text 'SAPL' is overlaid in red on the right side of the image.

SAPL



SAPL




Crest \$

S de Lannes



K pulmonaire





K pulmonaire



Hypertension art pulm



Osler



M Charcot

A close-up photograph of a person's hand and forearm. The skin is covered with a widespread, red, and scaly rash. The rash consists of numerous small, raised, red papules and larger, more confluent areas of redness with fine scaling. The distribution is most prominent on the back of the hand and the dorsal aspect of the forearm, extending towards the wrist. The fingers also show some involvement, with redness and scaling visible on the knuckles and the sides of the fingers. The overall appearance is characteristic of a severe allergic reaction or a severe form of eczema.

Still





dyshidrose

Dyshidrose palmaire









Bilan d'un acrosyndrome

- Examen clinique

- Local
- Général

Notion de terrain auto-immun

- Examens para-cliniques :

- Capillaroscopie
- Auto-Ac
- Radiographie thoracique
- Cryoglobulinémie
- NF réticulocytes